arXiv:1607.02727v1 [math.ST] 10 Jul 2016

Integral form of the COM–Poisson normalization constant

Tibor K. Pogány

Faculty of Maritime Studies, University of Rijeka, 51000 Rijeka, Studentska 2, Croatia
and
Institute of Applied Mathematics, Óbuda University, 1034 Budapest, Bécsi út 96/b, Hungary

Abstract

In this brief note an integral expression is presented for the COM–Poisson renormalization constant $Z(\lambda, \nu)$ on the real axis.

Keywords: CoM–Poisson renormalization constant $Z(\lambda, \nu)$, Cahen integral, Dirichlet series, Inverse Gamma function
2010 MSC: 62M10, 40C10, 62M20

1. Introduction and Motivation

Consider a rv X having classical COM–Poisson distribution initiated by Conway–Maxwell [1962] in the following form:

$$\mathsf{P}(X = n) = \frac{\lambda^n}{Z(\lambda, \nu)\,(n!)^\nu}, \qquad \lambda, \nu > 0; \quad n \in \mathbb{N}_0,$$

$Z(\lambda, \nu)$ being the reciprocal of the normalizing constant.

To the best of my knowledge, the only existing integral expression for $Z(\lambda, \nu)$ was given by Schmueli *et al.* [2005, p. 141, Eq. (40)] for positive integer ν as a multiple integral

$$Z(\lambda, \nu) = \frac{1}{(2\pi)^{\nu-1}} \int_{-\pi}^{\pi} \cdots \int_{-\pi}^{\pi} \exp\left\{ \sum_{j=1}^{\nu-1} \exp\left\{ \mathrm{i} x_j \right\} \right.$$

Email address: poganj@pfri.hr (Tibor K. Pogány)
URL: http://www.pfri.hr/~poganj (Tibor K. Pogány)

$$+\,\lambda\exp\Big\{-\mathrm{i}\sum_{j=1}^{\nu-1}x_j\Big\}\Big\}\,\mathrm{d}x_1\cdots\mathrm{d}x_{\nu-1}\,.$$

In the same case with $\nu\geq 1$ assumed to be an integer, Nadarajah reported on the closed form expression Nadarajah [2009, p. 619, Eq. (6)]

$$Z(\lambda,\nu)={}_0F_\nu(-;1,\cdots,1;\lambda)\,,$$

where ${}_0F_\nu$ stands for the familiar generalized hypergeometric function with zero upper and ν lower parameters.

Another fashion related approximation and extension results regarding $Z(\lambda,\nu)$ are numerous, see for instance the recent articles Schmueli *et al.* [2005], Gillispie–Christopher [2015], Şimşek–Iyengar [2015] and the references therein.

2. The integral expression for $Z(\lambda,\nu)$

Here a single definite integral expression is established for $Z(\lambda,\nu)$, for all positive ν.

Theorem 1. *For all $\lambda,\nu>0$ we have*

$$Z(\lambda,\nu)=\frac{1}{1-\lambda}+\frac{\nu}{\lambda-1}\int_1^\infty x^{-(\nu+1)}\,\lambda^{\left[\Gamma^{-1}(x)\right]}\,\mathrm{d}x\,,$$

where Γ^{-1} stands for the increasing branch of the inverse Gamma in the right half-plane, while $[x]$ denotes the integer part of some real x.

Proof. It turns out that

$$Z(\lambda,\nu)=\sum_{n\geq 0}\frac{\lambda^n}{(n!)^\nu}=\sum_{n\geq 0}\lambda^n\,\mathrm{e}^{-\nu\ln\Gamma(n+1)}\,,$$

is a classical Dirichlet series. The main tool we refer to is the Cahen formula for the Laplace integral representation of Dirichlet series (reported firstly without proof by Cahen [1894] and proved by Perron [1908]). Namely, the Dirichlet series

$$\mathscr{D}_{\boldsymbol{a}}(r)=\sum_{n\geq 1}a_n e^{-rb_n},$$

where $\Re(r) > 0$, possessing positive monotone increasing divergent to infinity sequence $(b_n)_{n\geq 1}$, has a Laplace integral representation Cahen [1894, p. 97]

$$\mathscr{D}_{\boldsymbol{a}}(r) = r \int_0^\infty e^{-rx} \sum_{n\colon\, b_n \leq x} a_n \,\mathrm{d}x = r \int_0^\infty e^{-rx} \sum_{n=1}^{[b^{-1}(x)]} a_n \,\mathrm{d}x\,,$$

since $b\colon \mathbb{R}_+ \mapsto \mathbb{R}_+$ is monotone, and there exists unique inverse b^{-1} for the function $b\colon \mathbb{R}_+ \mapsto \mathbb{R}_+$, being $b|_{\mathbb{N}} = (b_n)$, see also Pogány [2005].

Thus, bearing in mind that $b(x) \equiv \ln\Gamma(x+2)$ is monotone and invertible for $x \geq 1$, being $x > \alpha = 1.4616\cdots$ the abscissa of the minimum of the Gamma function Pedersen [2015], we only have to follow the previous derivation procedure:

$$\begin{aligned}
Z(\lambda,\nu) &= \sum_{n\geq 0} \lambda^n \,\mathrm{e}^{-\nu \ln\Gamma(n+1)} \\
&= 1 + \lambda + \lambda^2\nu \int_0^\infty \mathrm{e}^{-\nu x} \sum_{n\colon\, \ln\Gamma(n+2)\leq x} \lambda^{n-1} \,\mathrm{d}x \\
&= 1 + \lambda + \lambda^2\nu \int_0^\infty \mathrm{e}^{-\nu x} \sum_{n=1}^{\left[\Gamma^{-1}(\mathrm{e}^x)\right]-2} \lambda^{n-1} \,\mathrm{d}x \\
&= 1 + \lambda + \frac{\lambda^2\nu}{\lambda-1} \int_0^\infty \mathrm{e}^{-\nu x} \left(\lambda^{\left[\Gamma^{-1}(\mathrm{e}^x)\right]-2} - 1\right) \mathrm{d}x \\
&= \frac{1}{1-\lambda} + \frac{\nu}{\lambda-1} \int_1^\infty x^{-(\nu+1)} \,\lambda^{\left[\Gamma^{-1}(x)\right]} \,\mathrm{d}x\,,
\end{aligned}$$

which finishes the proof of the stated integral representation. □

Acknowledgement

I am grateful to Professor Satish Iyengar, University of Pittsburgh for discussion in which he introduced me the COM–Poisson distribution's open questions.

References

Cahen, E. (1894). Sur la fonction $\zeta(s)$ de Riemann et sur des fontions analogues. *Ann. Sci. l'École Norm. Sup. Sér. Math.* **11**, 75–164.

Conway, R. W. and Maxwell, W. L. (1962). A queuing model with state dependent service rates. *J. Industrial Engineering* **12**, 132–136.

Gillispie, S. B. and Christopher, G. C. (2015). Approximating the Conway-Maxwell-Poisson distribution normalization constant. *Statistics* **49**, No. 5, 1062–1073.

Nadarajah, S. (2009). Useful moment and CDF formulations for the COM-Poisson distribution. *Stat. Papers* **50**, 617–622.

Pedersen, H. L. (2015). Inverses of gamma functions. *Constr. Approx.* **41**, No. 2, 251–267.

Perron, O. X. (1908). Zur Theorie der Dirichletschen Reihen. *J. Reine Angew. Math.* **134**, 95–143.

Pogány, T. K. (2005). Integral representation of Mathieu $(\boldsymbol{a}, \boldsymbol{\lambda})$–series. *Integral Transforms Spec. Funct.* **16**, No.8, 685–689.

Shmueli, G., Minka, T. P., Kadane,J. B., Borle, S. and Boatwright, P. (2005). A useful distribution for fitting discrete data: revival of the Conway-Maxwell-Poisson distribution. *J. Roy. Statist. Soc. Ser. C* **54** No. 1, 127–142.

Şimşek, B. and S. Iyengar, S. (2015). Approximating the Conway–Maxwell–Poisson normalizing constant. *Filomat* (to appear).